Site_ID:	
Patient ID:	





- EPIS-AKI -

INTERNATIONAL OBSERVATIONAL STUDY TO EVALUATE THE EPIDEMIOLOGY OF SURGICAL-INDUCED ACUTE KIDNEY INJURY

CRF

INSTRUCTIONS FOR THE COMPLETION OF CASE REPORT FORMS

- Please use a black or dark-blue ballpoint pen to complete the case report form (CRF) and print legibly in English).
- In case of error, draw a single line through the incorrect entry, so the original entry is still visible and write the correct value next to it. Date and initial the correction.
- Dates must be entered in the order day / month / year. Write out the month using the first three letters

Example: April 24th 2020 must be <u>18/APR/2020</u>

In case of unknown day or month for a date, please indicate "UK"

Example: April 2020 must be $\underline{U}\underline{K}/\underline{A}\underline{P}\underline{R}/\underline{2}\underline{0}\underline{2}\underline{0}$

- ➤ Times must be entered as 24-hour clock time. Valid times are: <u>0 0</u>: <u>0 0</u> to <u>2 3</u>: 5 9. Must use leading zeros.
- ➤ If a portion of an examination / module was not performed for any reason or the data is unobtainable, please cross out the section concerned and state the reason using the following abbreviations as appropriate:

ND = not done **NA** = not applicable **UK** = unknown

Example: Weight _ U K . _

Please cross out unused repeating lines.

	e_ID:	EPIS-AKI Epidemiology of surgical induced acute kidney injury
INCL	USION CRITERIA	
The	following must be answered with YES otherwise the subject is not eligi	ble for the study
	Age ≥ 18	□₁yes □₀no
2.	Major surgeries with a duration of at least 2 h	
3.	Planned or unplanned admission to the ICU, IMC or PACU after	□₁yes □₀no
	surgery	
4.	Written informed consent	□₁yes □₀no
Exc	LUSION CRITERIA (inclusion only if none of the exclusion criteria are fulfilled)	
The	following must all answered with NO otherwise the subject is not eligil	ble for the study
1.	Pre-existing Acute Kidney Injury	□₁yes □₀no
2.	AKI within the last 3 months	□₁yes □₀no
3.	End-stage renal disease requiring Renal Replacement Therapy	□₁yes □₀no
4.	Kidney transplant	□₁yes □₀no
Nev	v Patient Entry	
INEV	VI AHLIVI LIVINI	
Adn	nission date hospital: / / [DD/MMM/YYYY]	

Admission date hospital:	/	_/	[DD/MMM/YYYY]
Inclusion date:	/	_/	[DD/MMM/YYYY]
Operation date:	/	/	[DD/MMM/YYYY]
Admission date ICU:	/	/	[DD/MMM/YYYY]

Site_ID:	



Patient ID:					induced acute kidney injury
PATIENT DATA					
Age	[years]				
Gender	□₁Male	\square_2 Female		□ ₉₉₉ Unkno	wn
Ethnicity	□₁Caucasian	□₂Black / A	frican descent	□₃Asian	□₄Hispanic
	□₅Other	Specify:			
Height	[cm]				
Weight	[kg]				
Creatinine baseline	□1	mg/dL	$\square_2\mu$ mol/L	□ ₉₉₉ Unkno	own
(known or presumed to have occurred within 7 days prior surgery):					

Site ID:		





MEDICAL HISTORY				
COMORBIDITIES				
Hypertension	□₁yes	□₀no		
Atrial fibrillation/flutter	□₁yes	□₀no		
Previous myocardial infarction	□₁yes	□₀no		
Congestive heart failure	□₁yes	□₀no		
If yes , please specify:	□¹NYHA I	□2NYHA II	□₃NYHA III	□ ₄ NYHA IV
Diabetes	□₁yes	□₀no		
If yes , please specify:	$\square_1 IDDM$	$\square_2 NIDDM$		
COPD	□₁yes	□₀no		
CKD [GFR < 60 mL/min]	□₁yes	□₀no		
If yes , please specify:	CKD stage:	□₃3 (eGFR 30	-59 ml/min)	
		□44 (eGFR 15	-29 ml/min)	
		□₅5 (eGFR < 1	L5 ml/min)	
Peripheral vascular disease	□₁yes	□₀no		
Previous stroke	□₁yes	□₀no		
ASA score:	□₁ASA 1	\square_2 ASA 2	□₃ASA 3	□₄ASA 4
MEDICATION	_	_		
Aspirin (ASS)	□₁yes	□₀no		
ACE inhibitors or ARBs	□₁yes	□₀no		
Beta-blockers	\square_1 yes	□₀no		
Diuretics	□₁yes	□₀no		
NSAIDs (except ASS)	□₁yes	□₀no		
Statins	□₁yes	□₀no		
Vasopressors	□₁yes	□₀no		
Use of contrast media one week prior to surgery	□₁yes	□₀no		

Site_ID:	FPI
Patient ID:	Epidemiolo induced acu
INTRAOPERATIVE DATA	
SURGICAL SPECIALTY (choose specialty responsible for surgical intervention)	
□₁Abdominal/General	
□₂Cardiac	
□ ₃ Gynaecologic/Obstetric	
□₄Neurosurgical	

If **yes**, please specify: _____

Type of surgery

 \square 5Orthopaedic

 \square ₆Thoracic

 \square_7 Trauma

 \square ₈Vascular

 \square 9Urologic

□10Others:

□1Open surgery	□₂Non-robotic minimal invasive surgery
□₃Robotic surgery	
$\square_{ exttt{4}}$ Planned combined n	ninimal invasive and open surgery
□₅Unplanned combine	d minimal invasive and open surgery

SURGICAL PROCEDURE

(See official website of International	l Classification of Health	Interventions: https://www.mitel.dimi.uniud.it.ichi):
 -	·_·	(Example: laparoscopic appendectomy code: KBO.JK.AB)
 -	·_·	Please add up to three codes if there was more than one
 -	··	surgical procedure performed at the same time.
D		

DURATION

Skin incision date/time:	///	::
	[DD/MMM/YYYY]	[HH:MM]
Skin suture date/time:	///	::
	[DD/MMM/YYYY]	[HH:MM]

Site_ID:	EPIS-AKI
Patient ID:	Epidemiology of surgical induced acute kidney injury
If CPB , please specify:	
Total CPB time: min	Total X-Clamp time: min
INTRAOPERATIVE DATA — CONTINUAT	ION (Data from start of anesthesia until end of anesthesia)
FLUID INTAKE	
Crystalloids:	[mL]
Colloids:	[mL]
Cell saver used:	□₁yes □₀no
(If a cellsaver is used please disregard the ce intake/output calculations.)	ell savers amount in the following and DO NOT include its volume in fluid
EC:	[mL]
TC:	[mL]
FFP:	[mL]
Оитрит	
Total blood loss:	[mL]
Urine catheter used:	□₁yes □₀no
Total urinary output:	[mL]
EPISODES OF HYPOTENSION (MAP < 55 mm)	Hg for more than 5 minutes): □₁yes □₀no
Application of vasopressors:	□₁yes □₀no
If yes , please specify:	□¹Norepinephrine □²Epinephrine □³Dobutamine □⁴Vasopressin □⁵Other:
Application of nephrotoxic agents:	□₁yes □₀no
If yes , please specify:	□₁ACEi/ARBs
	□ 2Aminoglycosides □ 3Amphotericin B
	□₄Cyclosporine/Tacrolimus

Site_ID:	EPIS—Z Epidemiology of s induced acute kid	urgical			
	Induced dedic Nu	icy injury			
	□₅NSAIDs (except Aspirin/ASS)				
	☐ Radiocontrast agents				
	□ ₇ Vancomycin				
	□₅Other:				
Application of diuretics:	□₁yes □₀no				
If yes , please specify:					
Intraoperative complications	□₁yes □₀no				
If yes , please specify:	□₁CPR				
	□ ₂ Arrhythmias				
	☐₃ Bleeding (Requiring at least one red blood cell unit)				
	\square ₄ Pulmonal complications (e.g., aspiration, brono	hospasm)			
	□ ₅ Allergic reaction				
	□₀Other:				

Site ID:	
_	

Patient ID:



POSTOPERATIVE DATA (from end of an	esthesia until end of postoperative c	lay 3)			
Start of Documentation date/time: (= end of anesthesia)	///	::			
(= end of anesthesia)	[DD/MMM/YYYY]	[HH:MM]			
End of Documentation date/time: (= end of postoperative day 3)		:: [HH:MM]			
APACHE II:	(Determined within 24 hours of admission to ICU)				
SAPS:	(Determined within 24 hours of admission to ICU)				
FLUID INTAKE (cumulative, i.v. and oral)					
Crystalloids:	_ [mL]				
Colloids:	[mL]				
EC:	_ [mL]				
TC:	_ [mL]				
FFP:	_ [mL]				
OUTPUT (cumulative)					
Total blood loss:	_ [mL] (including drainages)				
Total urinary output:	_ [mL]				
Urine catheter removed:	□₀Immediately post-ope	erative			
	□₁Day 1 □₂Day 2 □₃Da	ay 3			
	\square 4Removed later than d	ay 3			
DRUG APPLICATION					
Application of vasopressors:	□₁yes □₀no				
If yes , please specify:	: □₁Norepinephrine				
	□₂Epinephrine				
	□₃Dobutamine —				
	□₄Vasopressin				
	□₅Other:				

Site_ID	:	_					S-AKI
Patient	ID:	_					gy of surgical ute kidney injury
Applicat	ion of nephrotoxic a If yes , please	•	□2Am □3Am □4Cyc □5NSc □6Rac □7Van	Ei/ARBs ninoglycos nphoterici closporine AIDs (exc	sides in B e/Tacrolimus ept Aspirin/A st agents	SS)	
Applicat	ion of diuretics:		□₁yes	;	□₀no		
			□₀Pos	st-OP	□₁Day 1	□₂Day 2	□₃Day 3
Postopi	ERATIVE DATA CONTI	NUED (from 6	end of an	ıesthesia ur	ntil end of posto	perative day 3)	
POSTOPER	If yes , please		the □₂Ble □₃Re- □₄Pne □₅Sep	modynan rapy or incre eding (req operation eumonia osis	nic instability Pease of norepinep Juiring at least one	(requiring new ons hrine/epinephrine i e red blood cell unit	≥ 0.02µg/kg/min)
	IGO criteria for diagnosis o	of AKI					
Stage	Serum creatinine	1/12 -			Urine output		
1	≥ 0.3 mg/dL (≥ 26.5 µmol/L) increase in 48 h or 1.5–1.9-times baseline within the last 7 days		< 0.5 mL/kg/h f	or≥6 h			
2	2.0-2.9-times baseline		< 0.5 mL/kg/h f				
3	3-times baseline or $\geq 4.0 \text{ mg/dL}$ ($\geq 353.6 \mu \text{mol/L}$) increase or initiation of renal replacement therapy		< 0.3 mL/kg/h f anuria for ≥ 12				
Severity	dney injury (AKI): of AKI (worst case):	□₁yes □₀l	no L 🗆 2	KDIGO 2	□₃KDIGO 3	_	
Occurre	nce:	□ ₀ Post-OP		Day 1	□₂Day 2	□₃Day 3	
Duration	n of AKI:	□₁Transie	nt (< 48	Bh)	□₂Persiste	nt (> 48h)	

Site_ID:				EPIS-AKI Epidemiology of surgical induced acute kidney injury
Diagnosis (worst case AKI): □¹SCr	□₂Uı	rine output	□₃	Both
Bladder catheter removed		•	y 2 □₃	Day 3
Renal replacement therapy (RRT):	□₁yes	□₀no		
If yes , start of RRT:	/	/	[DD/MMN	M/YYYY]
Modality at time of in	nitiation:	□₁CRRT	□2 SLEDD	□₃IHD
Indication for initiation of RRT:	□₁Lung ede	ma	□₂Hyperv	olemia
	□₃Electrolyt	te derangement		
	□₄Uremia	□₅Anuria	□₀Others	

Site_ID: Patient ID:	EPIS-AKI Epidemiology of surgical induced acute kidney injury
Outcomes	
Date of ICU discharge:/ [DD/MMN	I/YYYY]
Date of Hospital discharge:/ [DD/MMN	I/YYYY]
Condition at hospital discharge: □₀Alive □₁Death	
Date of death:/[DD/MMM	I/YYYY]
Serum Creatinine at ICU discharge: \pi_1mg/dL \pi_2	μmol/L
at hospital discharge: □1mg/dL □2	μmol/L
RRT at hospital discharge: □¹Yes □₀No If yes : Total days RRT	
DAY 90 FOLLOW UP	
Index study date +90 Date of Follow Up:/(DD/MMM/YYYY)	
Condition: \square_0 Alive \square_1 Death Date of death : $____$	/[DD/MMM/YYYY]

Total days RRT (from hospital discharge until index study +90): _ _ _

□2μmol/l

Serum Creatinine: □1mg/dL

 \square_{999} unknown

 \square_{999} unknown